

11-1-2009

An Inductive Approach to Calculate the MLE for the Double Exponential Distribution

W.J. Hurley

Royal Military College of Canada, hurley-w@rmc.ca

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

Recommended Citation

Hurley, W. J. (2009) "An Inductive Approach to Calculate the MLE for the Double Exponential Distribution," *Journal of Modern Applied Statistical Methods*: Vol. 8 : Iss. 2 , Article 25.

DOI: 10.22237/jmasm/1257035040

An Inductive Approach to Calculate the MLE for the Double Exponential Distribution

W. J. Hurley
Royal Military College of Canada

Norton (1984) presented a calculation of the MLE for the parameter of the double exponential distribution based on the calculus. An inductive approach is presented here.

Key words: MLE, median, double exponential.

Introduction

Norton (1984) derived the MLE using a calculus argument. This article shows how to obtain it using a simple induction argument that depends only on knowing the shape of a function of sums of absolute values. Some introductory mathematical statistics textbooks, such as Hogg and Craig (1970) give the answer to be the median – although correct, this does not tell the whole story as Norton points out; this is emphasized here.

Methodology

It is useful to review the behavior of linear absolute value functions and sums of linear absolute value functions. For example, consider the function

$$g(x) = |1.8 - x|.$$

Its graph is shown in Figure 1. Note that it has a V-shape with a minimum at $x = 1.8$. Now consider a sum of two linear absolute value terms:

$$h(x) = |1.8 - x| + |3.2 - x|.$$

Plots of this function and its components, $|1.8 - x|$ and $|3.2 - x|$, are shown in Figure 2. Note that $h(x)$ takes a minimum at all points in the interval $1.8 \leq x \leq 3.2$.

The MLE

The double exponential distribution is given by

$$f(x) = \frac{1}{2} e^{-|x-\theta|}, \quad -\infty < x < \infty.$$

For the sample $\{x_1, x_2, \dots, x_n\}$, the log-likelihood function is

$$\ell(\theta) = n \ln(1/2) - \sum_i |x_i - \theta|.$$

Maximizing this function with respect to θ is equivalent to minimizing

$$g_n(\theta) = \sum_i |x_i - \theta|.$$

To obtain the MLE for general n , begin with the case $n = 1$ where $g_1(\theta) = |x_1 - \theta|$. This function has a minimum at $\theta = x_1$, hence, for $n = 1$, the MLE is

$$\theta^{MLE} = x_1.$$

W. J. Hurley is a Professor in the Department of Business Administration. Email: hurley-w@rmc.ca.

Figure 1: Plot of $g(x) = |x - 1.8|$





Figure 2: Plot of $h(x) = |1.8 - x| + |3.2 - x|$



INDUCTIVE MLE CALCULATION FOR THE DOUBLE EXPONENTIAL DISTRIBUTION

Now, consider the case $n = 2$. For the purposes herein it is useful to order the observations, thus, suppose that the sample is $\{x_{(1)}, x_{(2)}\}$ where $x_{(1)} < x_{(2)}$. The value of θ which minimizes must now be found using

$$g_2(\theta) = |x_{(1)} - \theta| + |x_{(2)} - \theta|.$$

Based on the above, this function takes the form

$$g_2(\theta) = \begin{cases} -2\theta + x_{(1)} + x_{(2)} & \theta \leq x_{(1)} \\ x_{(2)} - x_{(1)} & x_{(1)} \leq \theta \leq x_{(2)} \\ 2\theta - x_{(1)} - x_{(2)} & \theta \geq x_{(2)} \end{cases}$$

and has a minimum at any point θ in the interval $x_{(1)} \leq \theta \leq x_{(2)}$. Hence the MLE for $n = 2$ is

$$\theta^{MLE} = \lambda x_{(1)} + (1 - \lambda)x_{(2)}, \quad 0 \leq \lambda \leq 1.$$

For this case, the median is defined $(x_{(1)} + x_{(2)})/2$ and is a solution, but it is not the only solution.

Next, consider the case $n = 3$ with an ordered sample $x_{(1)} \leq x_{(2)} \leq x_{(3)}$. Using the

same graphical analysis, it can be shown that

$$g_3(\theta) = |x_{(1)} - \theta| + |x_{(2)} - \theta| + |x_{(3)} - \theta|$$

has a unique minimum at $\theta = x_{(2)}$, the median.

In the case $n = 4$, the solution is

$$\theta^{MLE} = \lambda x_{(2)} + (1 - \lambda)x_{(3)}, \quad 0 \leq \lambda \leq 1.$$

Thus, the median is a solution, but not the only solution.

Conclusion

Extending the argument for general n is straightforward. It is the median, $x_{((n+1)/2)}$, if n is odd and the generalized median, $\lambda x_{(n/2)} + (1 - \lambda)x_{(n/2+1)}$, when n is even.

References

Hogg, R. V., & Craig, A. T. (1970). *Introduction to Mathematical Statistics*, (3rd Ed.). New York, NY: MacMillan Publishing Company.

Norton, R. M. (1984). The double exponential distribution: Using calculus to find a maximum likelihood estimator. *The American Statistician*, 38(2), 135-136.