

11-2014

Reliability Estimates of Generalized Poisson Distribution and Generalized Geometric Series Distribution

Adil H. Khan

University of Kashmir, Srinagar, India, khanadil_192@yahoo.com

T R. Jan

University of Kashmir, Srinagar, India, drtrjan@gmail.com

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

Recommended Citation

Khan, Adil H. and Jan, T R. (2014) "Reliability Estimates of Generalized Poisson Distribution and Generalized Geometric Series Distribution," *Journal of Modern Applied Statistical Methods*: Vol. 13 : Iss. 2 , Article 20.
DOI: 10.22237/jmasm/1414815540

Reliability Estimates of Generalized Poisson Distribution and Generalized Geometric Series Distribution

Adil H. Khan

University of Kashmir
Srinagar, India

T. R. Jan

University of Kashmir
Srinagar, India

Discrete distributions have played an important role in the reliability theory. In order to obtain Bayes estimators, researchers have adopted various conventional techniques. Generalizing the results of Maiti (1995), Chaturvadi and Tomer (2002) dealt with the problem of estimating $P\{X_1, X_2, \dots, X_k \leq Y\}$, where random variables X and Y were assumed to follow a negative binomial distribution. Agit et al. obtained Bayesian estimates of the reliability functions and $P\{X_1, X_2, \dots, X_k \leq Y\}$ considering X and Y following binomial and Poisson distributions. The reliability function of the generalized Poisson and generalized geometric distribution is investigated. The expression for $P\{X_1, X_2, \dots, X_k \leq Y\}$ was obtained with X 's and Y following a Poisson distribution and some particular cases are shown.

Keywords: Generalized Poisson distribution generalized geometric distribution, reliability function, Bayes estimators

Introduction

Much research exists in the literature for estimating various parametric functions of several discrete distributions through classical and Bayesian approaches. Cacoullos and Charalambides (1975) obtained MVUE for truncated binomial and negative binomial distributions. Bayesian estimation of the parameter of binomial distribution has been considered by Chew (1971). Barton (1961) and Glasser (1962) obtained UMVUE of $P(X = x)$ for Poisson distribution. Blyth (1980) studied the absolute error of UMVUE of the probability of success of binomial distribution. For a random variable X following binomial distribution, Pulskanp (1990) has shown that the UMVUE of $P(X = x)$ is admissible under squared-error loss function

Adil H. Khan is a Research Scholar in the Post-Graduate Department of Statistics. Email him khanadil_192@yahoo.com. T. R. Jan is an Assistant Professor in the Post-Graduate Department of Statistics. Email him at drtrjan@gmail.com.

RELIABILITY ESTIMATES OF TWO GENERALIZED DISTRIBUTIONS

when $X = 0$ or n . Irony (1992) developed Bayesian estimation procedures related to Poisson distribution. Guttman (1958) and Patil (1963) provided UMVUEs of parametric functions of negative binomial distribution. Patil & Wani (1966) obtained UMVEUs of distribution function of various distributions. Roy & Mitra (1957) considered the problem of minimum variance unbiased estimation of univariate power series distribution. Patil (1978) generalized their results to multivariate modified power series distribution. Patil & Bildikar (1966) derived MVUE for logarithmic series distribution.

Discrete distributions have played important role in reliability theory. Kumar & Bhattacharya (1989) considered negative binomial distribution as the life time modal and obtained UMVUEs of the mean life and reliability function. Another measure of reliability is under stress-strength setup in the probability $Pr\{X \leq Y\}$, under the assumption that X and Y followed geometric distribution and derived UMVUE & Bayes estimator. Chaturvedi & Tomer (2002) considered classical & Bayesian Estimation procedures for the reliability function of the negative binomial distribution from a different approach generalizing the results of Maiti (1995), they dealt with the problem of estimating $P\{X_1, X_2, \dots, X_k \leq Y\}$, where random variables X & Y were assumed to follow negative binomial distributions. Chaturvadi, et al. (2007) considered Binomial and Poisson distribution and obtained the Bayesian estimators of reliability function and dealt with the problem of estimating $P\{X_1, X_2, \dots, X_k \leq Y\}$, where the random variables X & Y were assumed to follow binomial and Poisson distributions.

In order to obtain Bayes estimators of parameter and various parametric functions of different distributions, researchers have adopted a conventional technique, i.e., obtaining their posterior means. This article considers the Generalized Poisson and Generalized Geometric distributions and the problems of estimating reliability functions and $P = P\{X_1, X_2, \dots, X_k \leq Y\}$ from a Bayesian viewpoint. Bayes estimators of these parametric functions are derived. It is worth mentioning that in contrary to conventional approach, only estimators of factorial moments are needed to estimate these parametric functions and no separate dealing is needed.

Generalized Poisson Distribution

The random variable follows Generalized Poisson distribution with parameter λ and β if its pmf is

$$P(X = x) = \frac{\lambda^x (1 + x\beta)^{x-1} e^{-\lambda(\beta x + 1)}}{x!}, \quad x = 0, 1, 2, \dots, \lambda > 0, 0 < \beta < \frac{1}{\lambda} \quad (1)$$

The reliability function at a specific mission time, for example, $t_0 (\geq 0)$ is

$$R(t_0) = \sum_{x=t_0}^{\infty} \frac{\lambda^x (1 + x\beta)^{x-1} e^{-\lambda(\beta x + 1)}}{x!} \quad (2)$$

and the hazard rate function is

$$h(t_0) = \frac{P(t_0, \lambda, \beta)}{R(t_0)} = \left[\sum_{x=0}^{\infty} \frac{\lambda^x e^{-\lambda\beta x} (1 + (x + t_0)\beta)^{x+t_0-1}}{(1 + t_0\beta)^{t_0-1} (x + t_0)_{(x)}} \right]^{-1} \quad (3)$$

Let $\{X_i\}$, $i = 1, 2, 3, \dots, k$ be k independent random variables following a generalized Poisson distribution (1) with parameters λ_i and β (known) and Y is a random variable, independent of X 's following generalized Poisson distribution with parameter u . Denoting

$$X^* = \sum_{i=1}^k X_i \quad \text{and} \quad \lambda^* = \sum_{i=1}^k \lambda_i \quad (4)$$

From additive property of generalized Poisson distribution

$$P = \sum_{x^*=0}^{\infty} \sum_{y=x^*}^{\infty} p(x^*; \lambda^*) \cdot p(y; u) \quad (5)$$

Next, Bayes estimators of $R(t_0)$ and 'P' for generalized Poisson distribution are estimated.

Bayes Estimation Of $R(t_0)$ and 'P' For Generalized Poisson Distribution

The likelihood function given the random sample information $X = (X_1, X_2, \dots, X_n)$ is

$$L(\lambda, \beta / t) = \prod_{i=1}^n \left(\frac{(1 + x_i \beta)^{x_i - 1}}{x_i!} \right) \lambda^t e^{-\lambda(\beta t + n)} \tag{6}$$

where

$$t = \sum_{i=1}^n x_i$$

Because, $\lambda > 0$ consider the prior distribution for λ when β is known to be gamma with parameters (α, θ) and pdf

$$g(\lambda) = \frac{\theta^\alpha e^{-\theta \lambda} \lambda^{\alpha - 1}}{\Gamma(\alpha)}, \lambda, \alpha, \theta > 0 \tag{7}$$

From (6) and (7), the posterior density function of λ is given by

$$\prod(\lambda / t) = \frac{e^{-\lambda(\beta t + n + \theta)} \lambda^{t + \alpha - 1} (\beta t + n + \theta)^{t + \alpha}}{\Gamma(t + \alpha)} \tag{8}$$

The Bayesian estimator of λ^p , for $p > 0$, is given by

$$\begin{aligned} \hat{\lambda}^p &= \int_0^\infty \lambda^p \prod(\lambda / t) d\lambda \\ \hat{\lambda}^p &= \frac{\Gamma(t + \alpha + p)}{\Gamma(t + \alpha)} (\beta t + n + \theta)^{-p} \end{aligned} \tag{9}$$

Now, Equation (1) can be written as

$$P(x = x) = \frac{1}{x!} \sum_{r=0}^\infty (-1)^r \frac{(\beta x + 1)^{r + x - 1}}{r!} \lambda^{r + x}$$

On using (9) the Bayes estimator of $P(x;\lambda)$ at a specific point ‘X’ is

$$\begin{aligned} \hat{p}(x; \lambda) &= \frac{1}{x!} \sum_{r=0}^{\infty} (-1)^r \frac{(\beta x + 1)^{r+x-1}}{r!} \frac{\Gamma(t + \alpha + r + x)}{\Gamma(t + \alpha)} (\beta t + n + \theta)^{-(r+x)} \\ &= \frac{(\beta x + 1)^{x-1}}{(\beta t + n + \theta)^x} \cdot \frac{1}{x!} \sum_{r=0}^{\infty} (-1)^r \frac{(\beta x + 1)^r}{r!} \frac{(t + \alpha + r + x - 1)!}{(t + \alpha - 1)! (\beta t + n + \theta)^r} \quad (10) \\ &= (\beta x + 1)^{x-1} \binom{t + \alpha + x - 1}{x} \frac{(\beta t + n + \theta)^{t+\alpha}}{(\beta(t+x) + n + \theta + 1)^{t+\alpha+x}} \end{aligned}$$

Using (10) in (2), in order to obtain Bayesian estimator of $R(t_0)$, results in

$$\hat{R}(t_0) = \left(\frac{\beta t + n + \theta}{\beta(t+x) + n + \theta + 1} \right)^{t+\alpha} \sum_{x=t_0}^{\infty} \binom{t + \alpha + x - 1}{x} \frac{(\beta x + 1)^{x-1}}{(\beta(t+x) + n + \theta + 1)^x} \quad (11)$$

Also, for obtaining Bayesian estimator for ‘P’ we consider independent priors for λ^* and u to be gamma with parameters (α_1, θ_1) and (α_2, θ_2) respectively and using equations (4) and (10) is

$$\begin{aligned} &\sum_{x^*=0}^{\infty} \sum_{y=x^*}^{\infty} (\beta x^* + 1)^{x^*-1} (\beta y + 1)^{y-1} \binom{t_1 + \alpha_1 + x^* - 1}{x^*} \binom{t_2 + \alpha_2 + y - 1}{y} \\ \hat{P} &= \frac{\left(\beta t_1 + \sum_{i=1}^k n + \theta_1 \right)^{t_1 + \alpha_1} (\beta t_2 + m + \theta_2)^{t_2 + \alpha_2}}{\left(\beta(t_1 + x^*) + \sum_{i=1}^k n + \theta_1 + 1 \right)^{t_1 + \alpha_1 + x^*} (\beta(t_2 + y) + m + \theta_2 + 1)^{t_2 + \alpha_2 + y}} \quad (12) \end{aligned}$$

Special Cases

For $\beta = 0$, equation (1) reduces to Poisson distribution, therefore for $\beta = 0$, equations (11) and (12) give the Bayesian estimators for $R(t_0)$ and P (see Chaturvedi, et al., 2007) and are

$$\hat{R}_1(t_0) = \left(\frac{n + \theta}{n + \theta + 1}\right)^{t+\alpha} \sum_{x=t_0}^{\infty} \binom{t + \alpha + x - 1}{x} \frac{1}{(n + \theta + 1)^x}$$

$$\hat{P} = \frac{\sum_{x^*=0}^{\infty} \sum_{y=x^*}^{\infty} \binom{t_1 + \alpha_1 + x^* - 1}{x^*} \binom{t_2 + \alpha_2 + y - 1}{y} \left(\sum_{i=1}^k n + \theta_1\right)^{t_1 + \alpha_1} (m + \theta_2)^{t_2 + \alpha_2}}{\left(\sum_{i=1}^k n + \theta_1 + 1\right)^{t_1 + \alpha_1 + x^*} (m + \theta_2 + 1)^{t_2 + \alpha_2 + y}}$$

Generalized Geometric Distribution

The random variable ‘X’ follows Generalized Geometric distribution with parameters α and β if its pmf is

$$P(x = x) = \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \alpha^x (1 - \alpha)^{1 + (\beta - 1)x} \tag{13}$$

$0 < \alpha < 1, |\alpha\beta| < 1, x = 0, 1, 2, \dots$

The reliability function at a specific mission time, for example, $t_0 (\geq 0)$ is

$$R(t_0) = \sum_{x=t_0}^{\infty} \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \alpha^x (1 - \alpha)^{1 + (\beta - 1)x} \tag{14}$$

and the hazard rate function is

$$h(t_0) = \frac{P(t_0, \alpha, \beta)}{R(t_0)}$$

Bayes Estimation Of $R(t_0)$ For Generalized Geometric Distribution

The likelihood function given the random sample information $X = (X_1, X_2, \dots, X_n)$ is

$$L(\alpha, \beta / t) = \prod_{i=1}^n \left(\frac{1}{\beta x_i + 1} \binom{\beta x_i + 1}{x_i} \right) \alpha^t (1 - \alpha)^{n + (\beta - 1)t} \quad (15)$$

where

$$t = \sum_{i=1}^n x_i$$

Because, $0 < \alpha < 1$, it is assumed that the prior information about α when β is known from Beta distribution with pdf

$$g(\alpha) = \frac{\alpha^{a-1} (1 - \alpha)^{b-1}}{B(a, b)}, 0 < \alpha < 1, a > 0, b > 0 \quad (16)$$

The posterior distribution from (15) and (16) can be written as

$$\prod(\alpha / t) = \frac{\alpha^{t+a-1} (1 - \alpha)^{n + (\beta - 1)t + b - 1}}{B(t + a, n + (\beta - 1)t + b)} \quad (17)$$

RELIABILITY ESTIMATES OF TWO GENERALIZED DISTRIBUTIONS

The Bayesian estimator of α^p , for $p > 0$, is given by

$$\begin{aligned}\hat{\alpha}^p &= \int_0^{\infty} \alpha^p \prod(\alpha/t) d\alpha \\ \hat{\alpha}^p &= \frac{B(t+a+p, n+(\beta-1)t+b)}{B(t+a, n+(\beta-1)t+b)}\end{aligned}\tag{18}$$

Now, equation (13) can be written as

$$P(x=x) = \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \sum_{r=0}^{\infty} (-1)^r \binom{\beta x + 1 - x}{r} \alpha^{x+r}$$

On using (18) the Bayes estimator of $P(x;\lambda)$ at a specific point 'X' is

$$\begin{aligned}\hat{P}(x;\alpha) &= \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \sum_{r=0}^{\infty} (-1)^r \binom{\beta x + 1 - x}{r} \\ &\times \frac{B(t+a+x+r, n+(\beta-1)t+b)}{B(t+a, n+(\beta-1)t+b)} \\ &= \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \frac{B(t+a+x, n+1+(\beta-1)t+b)}{B(t+a, n+1+(\beta-1)t+b)}\end{aligned}\tag{19}$$

Using (19) in (14), in order to obtain Bayesian estimator of $R(t_0)$, results in

$$\hat{R}(t_0) = \sum_{x=t_0}^{\infty} \frac{1}{\beta x + 1} \binom{\beta x + 1}{x} \frac{B(t+a+x, n+1+(\beta-1)t+b)}{B(t+a, n+1+(\beta-1)t+b)}.$$

References

- Barton, D. E. (1961). Unbiased estimation of a set of probability. *Biometrika*, 48, 227-229.
- Blyth, C. R. (1980). Expected absolute error of the usual estimator of the binomial parameter. *American Statistician*, 34, 155-157.
- Cacoullos, T. & Charalambides, C. (1975). On minimum variance unbiased estimation for truncated binomial and negative binomial distributions. *Annals of the Institute of Statistical Mathematics*, 27, 235-244.
- Chaturvadi, A., Tiwari, N., & Kumar, S. (2007). Some remarks on classical and Bayesian reliability estimation of binomial and Poisson distributions. *Statistical Papers*, 48, 683-693.
- Chaturvedi, A. & Tomer, S. K. (2002). Classical and Bayesian Reliability estimation of the negative binomial distribution. *Journal of Applied Statistical Science*, 11(1), 33-43.
- Chew, V. (1971). Point estimation of the parameter of the binomial distribution. *American Statistician*, 25, 47-50.
- Glasser, G. J. (1962). Minimum variance unbiased estimators for Poisson probabilities. *Technometrics*, 4, 409-418.
- Guttman, I. (1958). A note on series solution of a problem in estimation. *Biometrika*, 45, 565-567.
- Irony, T. Z. (1992). Bayesian estimation for discrete distribution. *Journal of Applied Statistics*, 19, 38-47.
- Kumar, S. & Bhattacharaya, S. K. (1989). Reliability estimation for negative binomial distribution. *Assam Statistical Review*, 3, 104-107.
- Maiti, S. S. (1995). Estimation of $Pr\{X \leq Y\}$ in the Geometric case. *Journal of the Indian Statistical Association*, 33, 87-91.
- Patil, G. P. (1963). Minimum Variance unbiased estimation of generalized Poisson distribution and certain problems of additive number theory. *Annals of Mathematical Statistics*, 34, 1050-1056.
- Patil, G. P. & Bildikar, S. (1966). On minimum variance unbiased estimation for the logarithmic series distribution. *Sankhya*, A28, 239-250.
- Patil, G. P. & Wani, J. K. (1966). Minimum variance unbiased estimation of the discretion function admitting a sufficient statistics. *Annals of Mathematical Statistics*, 18, 39-47.

RELIABILITY ESTIMATES OF TWO GENERALIZED DISTRIBUTIONS

Patil, S. R. (1978). Minimum variance unbiased estimation of multivariate modified power series distribution. *Metrika*, 25, 155-161.

Pulskamp, R. (1990). A note on the estimation of binomial probabilities. *American Statisticians*, 44, 293-295.

Roy, J. & Mitra, S. K. (1957). Unbiased minimum variance estimation in a class of discrete distributions. *Sankhya*, 18, 371-378.